

Quantum interference in bosonic and fermionic matter-wave amplification

H. Deng¹ and Y. Yamamoto^{1,2}

¹*Quantum Entanglement Project, ICORP, JST
Edward L. Ginzton Laboratory,
Stanford University, Stanford, CA 94305, USA*

²*NTT Basic Research Laboratories,
Atsugi-shi, Kanagawa, Japan*

We investigate the quantum interference effects in two types of matter-wave mixing experiments: one with initial matter waves prepared in independent Fock states (type I) and the other with each individual particle prepared in a same coherent superposition of states (type II). In the type I experiment, a symmetric wavefunction of bosons leads to constructive quantum interference and shows final state stimulation, while an anti-symmetric wavefunction of fermions results in destructive quantum interference and inhibited matter wave mixing. In the type II experiment, a coherent superposition state leads to constructive quantum interference and enhanced matter wave mixing for both bosons and fermions, independent of their quantum statistics.

With the realization of Bose-Einstein condensation (BEC) in atoms, bosonic final state stimulation involving atom condensates has been studied in superradiance of atoms [1], four-wave mixing (FWM)[2], and matter wave amplification[3, 4]. Following these work, it was pointed out that these phenomena are not unique in boson systems but also possible in fermion systems[5, 6]. In these experiments ([1]-[4]), the input matter waves, characterized by their momenta, are all prepared from a same condensate by a coherent partition process of each individual particle. Hence the observed nonlinearity can be understood as collective enhancement effect, analogous to Dicke super-radiance in an ensemble of two-level atoms, and does not depend on the quantum statistics of the particles. We call such experiments *type II* in this paper. In another kind of experiment, which we call *type I* here, all input matter waves consist of independent real populations of the particle. In this case, final state stimulation occurs in a boson system, while inhibition of matter-wave mixing is expected in a fermion system. Type I experiment has not yet been performed with atomic BECs, but has been recently demonstrated with exciton-polaritons in semiconductors[7, 8].

We investigate the two types of matter-wave mixing in terms of the quantum interference among different paths, naturally originating from the symmetrization (anti-symmetrization) procedure for type I, or artificially created from the coherent superposition state for type II. Figure 1 shows a model FWM experiment where two input states $|\phi\rangle$ and $|\psi\rangle$ elastically scatter into two output states $|u\rangle$ and $|v\rangle$ via the two possible processes (A) and (B), with scattering amplitudes S_A and S_B , respectively[9]. The scattering processes are governed by a unitary operator $\hat{U}_{int}$, s.t.,

$$S_A = {}_2\langle u|_1\langle v|\hat{U}_{int}(t)|\phi\rangle_1|\psi\rangle_2 = {}_2\langle v|_1\langle u|\hat{U}_{int}(t)|\psi\rangle_1|\phi\rangle_2, \quad (1)$$

$$S_B = {}_2\langle v|_1\langle u|\hat{U}_{int}(t)|\phi\rangle_1|\psi\rangle_2 = {}_2\langle u|_1\langle v|\hat{U}_{int}(t)|\psi\rangle_1|\phi\rangle_2. \quad (2)$$

Type I experiment In type I experiment, the four initial matter waves have definite populations (Fock states). Let's first consider the case where there is one particle in each of the two input states and one particle in one output state, with the initial state of the system:

$$\begin{aligned} |i_I\rangle &= \sum_{i \neq j} \hat{P}_{ij} [|\phi\rangle_1 |v\rangle_2 |\psi\rangle_3] \\ &= \frac{1}{\sqrt{6}} [|\phi\rangle_1 |v\rangle_2 |\psi\rangle_3 \pm |v\rangle_1 |\phi\rangle_2 |\psi\rangle_3 \pm |\psi\rangle_1 |v\rangle_2 |\phi\rangle_3 \\ &\quad + |v\rangle_1 |\psi\rangle_2 |\phi\rangle_3 \pm |\phi\rangle_1 |\psi\rangle_2 |v\rangle_3 + |\psi\rangle_1 |\phi\rangle_2 |v\rangle_3], \quad (3) \end{aligned}$$

where $\hat{P}_{ij}$ is the symmetrization or anti-symmetrization operators[9]. The upper sign is for bosons, and the lower sign is for fermions, in accordance with symmetrization




FIG. 1:
Illustration of the two scattering processes (A) and (B) in FWM experiments.




FIG. 2:

Possible paths of scattering leading to the final state (a) $|v_1|v_2|u_3$ in type I experiment, (b) $|v_1|v_2|u_3$ in type II experiment, and (c) $|\phi_1|v_2|u_3$ in type II experiment. In (a), the upper sign is for bosons, the lower sign is for fermions. The total scattering amplitudes is enhanced for bosons, and is suppressed to zero for fermions in this type I experiment. In (b) and (c), the scattering amplitudes are the same for both bosons and fermions. It is enhanced in (c), but there is no interference leading to enhanced or suppressed scattering in (c).

postulate for bosons and fermions, respectively. Scattering results in final states with two particles in state $|v\rangle$ and one particle in state $|u\rangle$. Take a final state $|v_1|v_2|u_3\rangle$ as an example (Fig. 2(a)), it can be reached by scattering (A) or (B) from each of the first four terms in the initial state $|i_I\rangle$. The corresponding scattering amplitudes for the four paths are S_AC , $\pm S_AC$, $\pm S_BC$, and

S_BC , where C is a real normalization factor, $C = 1/\sqrt{6}$ in this example. Thus the total scattering amplitude for bosons adds up to $2(S_A + S_B)C$ as a result of constructive quantum interference; while for fermions it is suppressed to zero due to destructive quantum interference. This illustrates how quantum interference leads to final state stimulation for bosons, and inhibited FWM for fermions.

In general, if the initial matter waves of a boson system consists of n_1 particles in $|\phi\rangle$, n_2 particles in $|\psi\rangle$, n_3 particles in $|v\rangle$, and $n = n_1 + n_2 + n_3$ particles in total, the initial-state is:

$$|i\rangle_{IB} = \frac{1}{\sqrt{N}} \left[\prod_{i=1}^{n_1} |\phi\rangle_i \prod_{j=n_1+1}^{n_1+n_2} |\psi\rangle_j \prod_{k=n_1+n_2+1}^n |v\rangle_k + \text{permutation terms due to symmetrization postulate} \right]. \quad (4)$$

All the permutations add to a total number of $N = \binom{n}{n_1} \binom{n-n_1}{n_2}$ different terms in the bracket. Any one of the n_1 particles in $|\phi\rangle$ and any one of the n_2 particles in $|\psi\rangle$ can scatter into $|v\rangle$ and $|u\rangle$ via the two processes (A) and (B). Process (A) results in a total of $N_2 = N \binom{n_1}{1} \binom{n_2}{1}$ terms in the final state $|\tilde{f}\rangle_{IB} = \hat{U}_{int}|i\rangle_{IB}$. Each of these terms has one particle in $|u\rangle$, $n_1 - 1$ particles in $|\phi\rangle$, $n_2 - 1$ particles in $|\psi\rangle$, and $n_3 + 1$ particles in $|v\rangle$. Hence there are only $N_3 = \binom{n}{1} \binom{n-1}{n_1-1} \binom{n-n_1}{n_2-1}$ physically distinct terms in the final state. Due to the symmetry property of the initial state (4), all these N_3 terms have the identical probability amplitude $S_{ACIB} = S_A(\frac{N_2}{N_3})/\sqrt{N}$. Same analysis applies to the process (B) except that the probability amplitude of each term is $S_{BCIB} = S_B(\frac{N_2}{N_3})/\sqrt{N}$.

The normalized final state is $|f\rangle_{IB} = |\tilde{f}\rangle_{IB} / \sqrt{IB\langle\tilde{f}|\tilde{f}\rangle_{IB}}$. The scattering amplitude is:

$$\begin{aligned} a_{IB} &= IB\langle f|\hat{U}_{int}|i\rangle_{IB} = IB\langle f|\tilde{f}\rangle_{IB} = \sqrt{IB\langle\tilde{f}|\tilde{f}\rangle_{IB}} \\ &= \sqrt{N_3 c_{IB}^2} |S_A + S_B| \\ &= \sqrt{n_1 n_2 (n_3 + 1)} |S_A + S_B|. \end{aligned} \quad (5)$$

It shows that the scattering rate is proportional to the product of the numbers of particles (n_1 and n_2) in the input states and is enhanced by the initial occupancy n_3 of the output state $|v\rangle$.

In the case of a fermion system, the initial state is:

$$|i\rangle_{IF} = \frac{1}{\sqrt{n!}} [|\phi^{(1)}\rangle_1 |\phi^{(2)}\rangle_2 \dots |\phi^{(n_1)}\rangle_{n_1} |\psi^{(1)}\rangle_{n_1+1} \dots |\psi^{(n_2)}\rangle_{n_1+n_2} |v^{(1)}\rangle_{n_1+n_2+1} \dots |v^{(n_3)}\rangle_n + \text{permutation terms due to anti-symmetrization postulate}], \quad (6)$$

Here the superscripts (1), (2), ... are labels of another

quantum number q , which is conserved during the scat-

tering processes. Thus, for example, the n_1 particles occupy nearly degenerate but distinct states $|\phi^{(1)}\rangle_1, |\phi^{(2)}\rangle_2$, etc. There are a total of $n!$ physically distinct terms in the bracket. Due to the anti-symmetrized form of the initial state, scattering amplitude cancel out exactly between different paths if they lead to a final state with two

particles in a same state $|v\rangle$. For a scattering process in which two particles are scattered into initially unoccupied $|v\rangle$ and $|u\rangle$, there is no other paths interfering with it. Explicitly, the total scattering amplitude ${}_IF\langle f|\hat{U}_{int}|i\rangle_{IF}$ is:

$$a_{IF} = \begin{cases} 0, & \text{if } n_3 \geq n_1, n_2, \\ \sqrt{(n_1 - n_3)n_2}|S_A|, & \text{if } n_1 > n_3 \geq n_2, \\ \sqrt{(n_2 - n_3)n_1}|S_B|, & \text{if } n_2 > n_3 \geq n_1, \\ [(n_1 - n_3)n_2|S_A|^2 + (n_2 - n_3)n_1|S_B|^2 + 2(n_2 - n_3)(S_A S_B^* + S_A^* S_B)]^{1/2}, & \text{if } n_1 > n_2 > n_3, \\ [(n_1 - n_3)n_2|S_A|^2 + (n_2 - n_3)n_1|S_B|^2 + 2(n_1 - n_3)(S_A S_B^* + S_A^* S_B)]^{1/2}, & \text{if } n_2 > n_1 > n_3. \end{cases} \quad (7)$$

It shows that if there are initially more particles in an ensemble of nearly degenerate states $|v\rangle$ than in $|\phi\rangle$ and $|\psi\rangle$, the scattering into $|v\rangle$ and $|u\rangle$ is completely suppressed. Otherwise, the amplitude is non-zero but still suppressed by the increase of n_3 .

Type II experiment In contrast to the type I experiment, each particle in the initial matter waves of a type II experiment is in a same coherent superposition of the states $|\phi\rangle, |\psi\rangle$, and $|v\rangle$. In a boson system, if there are n particles in total, and each particle prepared in an identical superposition state, the initial-state wavefunction of bosons is:

$$|i\rangle_{IIB} = \prod_{i=1}^n \left(\sqrt{\frac{1-\epsilon}{2}} |\phi\rangle_i + \sqrt{\frac{1-\epsilon}{2}} |\psi\rangle_i + \sqrt{\epsilon} |v\rangle_i \right) \quad (8)$$

The expansion of (8) consists of a total of 3^n different terms. There are $N_{mk} = \binom{n}{m} \binom{n-m}{k}$ terms which have m particles in $|\phi\rangle$, k particles in $|\psi\rangle$ and $(n-m-k)$ particles in $|v\rangle$. Here m takes values from 0 to n , for each m , k takes values from 0 to $n-m$. We call this group of N_{mk} terms as (m,k) group. All terms in the same (m,k) group have the same probability amplitude $c_{mk}^0 = (\frac{1-\epsilon}{2})^{n/2} \eta^{\frac{n-m-k}{2}}$, where $\eta = \frac{2\epsilon}{1-\epsilon}$. It is obvious from the expansion that $|i\rangle_{IIB}$ is already fully symmetric, and no additional symmetrization procedure is necessary.

For each (m,k) group, possible scattering of a pair of $|\phi\rangle_i$ and $|\psi\rangle_j$ into $|v\rangle_i |u\rangle_j$ via process (A) results in a total of $N'_2 = mkN_{mk}$ terms, each of which has one particle in $|u\rangle$, $(m-1)$ particles in $|\phi\rangle$, $(k-1)$ particles in $|\psi\rangle$

and $(n-m-k+1)$ particles in $|v\rangle$. However, there are only $N'_3 = \binom{n}{1} \binom{n-1}{m-1} \binom{n-m}{k-1}$ physically distinct terms. Since the initial state is symmetric, the initial group (m,k) is scattered into N'_3 different terms, all with the same probability amplitude $S_A c'_{mk} = S_A c_{mk}^0 \frac{N'_3}{N_3} = c_{mk}^0 \cdot 2(n-m-k+1)$. Similarly, scattering via process (B) contributes $S_B c'_{mk}$ to the probability amplitude. The final state $|\tilde{f}\rangle_{IIB} = \hat{U}_{int}|i\rangle_{IIB}$ is a sum of terms scattered from all (m,k) groups. Hence the total scattering amplitude is:

$$\begin{aligned} a_{IIB} &= \sqrt{{}_{IIB}\langle \tilde{f} | \tilde{f} \rangle_{IIB}} \\ &= \sqrt{\sum_{m=1}^{n-1} \sum_{k=1}^{n-m} N'_3(m,k) (c'_{mk} |S_A + S_B|)^2} \\ &= \sqrt{\left[\frac{1-\epsilon}{2} n \right] \left[\frac{1-\epsilon}{2} (n-1) \right] [\epsilon(n-2) + 1] |S_A + S_B|}. \end{aligned} \quad (9)$$

When $n \gg 1$, the scattering rate is again proportional to the product of the average numbers of particles in the two input states (the first two terms in the last line of (9) corresponding to n_1 and n_2 in type I experiment), and is enhanced by the final state population by a factor $\epsilon(n-2) + 1$, where $\epsilon(n-2)$, the average population in $|v\rangle$, corresponds to n_3 in the type I experiment.

For type II experiment, a fermion system has exactly the same scattering amplitude. The initial state for fermions is

$$\begin{aligned} |i\rangle_{IIF} &= \frac{1}{N'} \left[\left(\frac{1-\epsilon}{2} \right)^{n/2} \prod_{i=1}^n (|\phi^{(i)}\rangle_i + |\psi^{(i)}\rangle_i + \sqrt{\eta} |v^{(i)}\rangle_i) \right. \\ &\quad \left. + \text{permutation terms due to anti-symmetrization postulate} \right], \end{aligned} \quad (10)$$

where N' is a normalization factor. Since each particle has a different quantum number q , and q is conserved under the operation of $\hat{U}_{int}$, terms in the expansions of different anti-symmetrization groups do not interfere with each other, even after scattering. At the same time, all anti-symmetrization groups have identical scattering characteristics. Therefore it is sufficient to consider only the first line of (10), setting $N' = 1$. In another word, symmetrization postulate and thus quantum statistics does not affect the scattering amplitude in type II experiment. Moreover, the label for quantum number q in (10) has a one to one correspondence to the label of the particle number, so the label for quantum number q can be suppress and (10) is reduced to the same form as (8), leading to $a_{IIF} = a_{IIB}$.

We again consider a simple case of three particles, each particle occupying the three states $|\phi\rangle$, $|\psi\rangle$ and $|v\rangle$ with equal probability. Then the initial state of bosons and the reduced initial state of fermions has the same form:

$$|i\rangle = (|\phi\rangle_1 + |\psi\rangle_1 + |v\rangle_1) \otimes (|\phi\rangle_2 + |\psi\rangle_2 + |v\rangle_2) \\ \otimes (|\phi\rangle_3 + |\psi\rangle_3 + |v\rangle_3)$$

To reach a final state $|v\rangle_1|v\rangle_2|u\rangle_3$, there are four possible paths, as illustrated in Fig. 2(b). The corresponding four terms in the initial state $|i\rangle$ originate from a coherent superposition state instead of the symmetrization or anti-symmetrization procedure. Therefore paths from each process are additive for both bosons and fermions. The constructive interference between different paths leads to enhanced scattering amplitude. To have an intuitive picture of the enhancement by the final state occupancy (corresponding to the $\epsilon(n-2)$ term in (9)), we consider the case where there are no particle in $|v\rangle$ before scattering. Then only two paths are possible, as shown in Fig. 2(c), and no interference terms to lead to enhancement in this case.

As discussed above, type II experiment will produce identical enhancement in scattering amplitude for both boson and fermion systems, given that the initial state

of the system is prepared as a coherent superposition of all three states $|\phi\rangle$, $|\psi\rangle$ and $|v\rangle$. The enhancement comes from constructive multi-particle interference, where the different paths are created by preparing the initial state in a coherent superposition-state. Type I experiment, however, will reveal final state stimulation for bosons and inhibited-FWM for fermions (Pauli blocking). The enhancement and inhibition in this case come from constructive and destructive multi-particle interference, where the different paths stem from the symmetrization and anti-symmetrization postulate. So only type I experiment tests the true quantum statistics of the system. The final state stimulation[7], matter wave amplification[8] and condensation of exciton polaritons[10] have been demonstrated in this type of experiment, but the counterpart experiments in atomic systems are yet to be observed.

-
- [1] S. Inouye, A. Chikkatur, D. Stamper-Kurn, J. Stenger, D. Pritchard, and W. Ketterle, *Science* **285**, 571 (1999).
 - [2] L. Deng, E. Hagley, J. Wen, M. Trippenbach, Y. Band, P. Julienne, J. Simsarian, K. Helmerson, S. Rolston, and W. Phillips, *Nature* **398**, 218 (1999).
 - [3] S. Inouye, T. Pfau, S. Gupta, A. Chikkatur, A. Gorlitz, D. Pritchard, and W. Ketterle, *Nature* **402**, 641 (1999).
 - [4] M. Kozuma, Y. Suzuki, Y. Torii, T. Sugiura, T. Kuga, E. Hagley, and L. Deng, *Science* **286**, 2309 (1999).
 - [5] M. Moore and P. Meystre, *Phys. Rev. Lett.* **86**, 4199 (2001).
 - [6] W. Ketterle and S. Inouye, *Phys. Rev. Lett.* **86**, 4203 (2001).
 - [7] R. Huang, F. Tassone, and Y. Yamamoto, *Phys. Rev. B* **61**, R7854 (2000).
 - [8] R. Huang, Y. Yamamoto, R. André, J. Bleuse, M. Muller, and H. Ulmer-Tuffigo, *Phys. Rev. B* **65**, 165314 (2002).
 - [9] J. J. Sakurai, *Modern Quantum Mechanics* (Addison-Wesley, New York, 1995).
 - [10] H. Deng, G. Weihs, C. Santori, J. Bloch, and Y. Yamamoto, *Science* **298**, 199 (2002).